



## Redox reactions of cobalt (III) Complexes of $\alpha$ -hydroxy acids by quinolinium chloro chromate (QCC) in Surfactants

\* K Murali, M Shreelekha, B Divya

Department of Chemistry, Vivekanandha College of Arts & Sciences for women, Sankagiri, Tamil Nadu, India

### Abstract

The kinetics of one electron transfer mode seems to be unavailable for QCC with Cobalt (III) bound and unbound complexes of  $\alpha$ -hydroxy acids in surfactant medium, QCC oxidizes Cobalt (III) bound and unbound  $\alpha$ -hydroxyl acids. It rules out the synchronous C-C bond fission and electron transfer to Cobalt (III) centre. Oxidation of above complexes increases with increase in temperature. The increase in the rate is observed with increase in the concentration of the surfactant. The added DDAC enhances the rate of oxidation of a reaction much more than Polysorbate 80. Similar trends has been observed in lactato, glycolato and Mandelato Co (III) complexes.

**Keywords:** quinolinium chloro chromate (QCC), polysorbate 80, dimethyl diocta decyl ammonium chloride (DDAC)

### 1. Introduction

Evolution 60 Thermo spectrophotometer has been employed to study the oxidation of  $\alpha$ -hydroxy acids such as mandelic acid, lactic acid, glycolic acid and their Cobalt (III) Complexes using Quinolinium Chloro Chromate as an oxidant in the presence of surfactant. One equivalent oxidant like Ce (IV) induced electron transfer in pentaamminecobalt (III) Complexes of  $\alpha$ -hydroxy acids results in nearly 100% reduction at Cobalt (III) centre with synchronous C-C bond fission and decarboxylation. Such an electron transfer route seems to be unavailable for Quinolinium Chloro Chromate in its reaction with cobalt (III) bound and unbound  $\alpha$ -hydroxy acids in surfactant medium. Quinolinium Chloro Chromate oxidizes cobalt (III) bound and unbound  $\alpha$ -hydroxy acids to respective keto acid cobalt (III) complexes in Polysorbate 80 and Dimethyl Diocta Decyl Ammonium Chloride (DDAC) possibly the transition state is more electron deficient. Such a transition state can be envisaged only when the C-H bond fission occurs in the slow step with a hydride ion transfer.

The absence of formation of cobalt (II) rules out the synchronous C-C bond fission and electron transfer to cobalt (III). The thermodynamic parameters are in consistent with bimolecular reaction. The rate of QCC oxidation of cobalt (III) Mandelato, Lactato and Glycolato complexes depends on the first power of QCC concentration. Similarly the reaction between QCC and unbound  $\alpha$ -hydroxy acid exhibits first order kinetics with respect to concentration of QCC. Of the three complexes lactate Cobalt (III) complexes react faster than mandelato and glycolato complexes, and similar trend is followed in the unbound ligands Polysorbate 80. Oxidation is an important process in organic chemistry and introduction of new economic and effective reagents for oxidation under mild and anhydrous conditions constitutes a standing challenge. QCC is an effective oxidant which is non-hygroscopic, non-photosensitive, stable yellow orange solid which is freely soluble in water, acetic acid, N,N-dimethyl formamide etc.

### 2. Materials and Methods

The surfactants used in the present work are Polysorbate 80 and Dimethyl Diocta Decyl Ammonium Chloride (DDAC). The surfactants are purified by adopting earlier procedure. The surfactants Polysorbate 80 and DDAC were purchased from (Sigma-Aldrich India 98%), mandelic, lactic and glycolic acids from (Karnataka Fine chemical. India 95%) Pentaamminecobalt (III) complexes of  $\alpha$ -hydroxy acids were prepared using 'Fan and Gould2, 10. Double distilled (deionised and CO<sub>2</sub> free) water was used as a solvent and HClO<sub>4</sub> (E. Merck India 95%) was standardized using standard Sodium carbonate (BDH, AR) solution with methyl orange as an indicator. For the QCC oxidation of Co (III) Complexes of  $\alpha$ -hydroxy acids and unbound ligands<sup>11, 12</sup>.


The rate measurement were made at  $34 \pm 0.2^\circ\text{C}$  in 100% aqueous medium and temperature was controlled by electrically operated thermostat. The total volume of reaction mixture in the spectrophotometric cell was kept as 2.5 ml in each kinetic run. An Evolution60 Thermo spectrophotometer fitted with recording and thermosetting arrangement was used to follow the rate of their action. Rate of this QCC oxidant with unbound ligand and Cobalt (III) bound complexes were calculated from observed decrease in absorbance at 400nm.

The excess of the reductant was used in kinetic runs. It gives pseudo first order rate constant. It was determined from the linear plot of the  $\ln A$  versus time. Reproducible result obtained giving good first order plot. The stoichiometric studies for the QCC oxidation of pentaamminecobalt (III) complexes of  $\alpha$ -hydroxy acid and unbound ligand in the presence of micelles were carried out at  $34 \pm 2^\circ\text{C}$ . It was observed that the cobalt (II) formation was negligibly small.

#### 2.1 Preparation of Quinolinium Chloro Chromate (QCC).

In to a 1-l beaker containing Hydrochloric acid (6 M, 184 ml) was added Chromium trioxide (100 g 1 mole) rapidly. The mixture was thoroughly stirred for five minutes and the resulting homogeneous solution was cooled to  $15^\circ\text{C}$ . To this

mixture quinoline (129.6 g 1 mole) was added drop wise over a period of half an hour. After the addition of quinoline was over the mixture was again cooled to 15° C. This resulted in precipitation of Quinolinium Chloro chromate in the form of a yellow powder. The powder was filtered off and dried under vacuum for three hours, melting point 239 °, 96 % yield.



### 3. Result and Discussion

Kinetic study of the oxidation of pentaamminecobalt (III) complexes of  $\alpha$ -hydroxy acid by QCC in surfactant medium dependence of rate on QCC concentration in bound ligand. The rate of oxidation of lactate cobalt (III) complexes depends on QCC concentration, the specific rate calculated remains constant (Table 1) and Graph of logarithm of QCC concentration versus time (Fig. 1) are linear.

From the slope of these graphs, the specific rate calculated agrees with those obtained from integrated rate equation suggesting first order dependence on QCC concentration.

When the concentration of QCC is varied from 1.00 to 5.00  $\times 10^{-3}$  mol dm<sup>-3</sup> at a fixed [Cobalt (III)] and [HClO<sub>4</sub>]. Specific rates remain constant. Then the of rate of disappearance of Cr (VI) is given by equation. (1)

$$-d [\text{Cr (VI)}] / dt = k [\text{Cr (VI)}] \quad \dots (1)$$

At a particular QCC concentration with increases in mandelato / lactato / glycolato cobalt (III) concentration in the range 1.00 to 4.00  $\times 10^{-3}$  mol dm<sup>-3</sup> there is a proportional increases in the rate of oxidation (Table 2). The slope of nearly unity is obtained from a linear graph of logarithm a (Fig. 2) of specific rate (k in s-1) versus logarithm of Co (III) concentration in each case suggesting first order rate dependence of rate on [Co (III)]. Hence the rate law for the Cr (VI) oxidation of cobalt (III) bound of  $\alpha$ -hydroxy acids is given by equation.(2).

$$-d [\text{Cr (VI)}] / dt = k_2 [\text{Cr (VI)}] [\text{Co (III)}] \quad \dots (2)$$

**Dependence of rate on QCC concentration in surfactant for Cobalt (III) complexes of  $\alpha$ - Hydroxyacid:** The rate of oxidation of lactato Cobalt (III) complexes depends on QCC concentration. In any specific run the change in concentration of QCC, the specific rate calculated remains constant (Table-3) and graphs of logarithm of QCC concentration versus time are linear. (Fig.3)

From the slope of these graphs, the specific rate calculated agrees with those obtained from integrated rate equation, suggesting first order dependence on QCC concentration.

When concentration of QCC is varied from 1.00 to 4.00  $\times 10^{-3}$  mol dm<sup>-3</sup> at a fixed [Co (III)] and [HClO<sub>4</sub>] specific rates remains constant. Then the rate of disappearance of Cr (VI) is given by equation 3.

$$-d [\text{Cr (VI)}] / dt = K_1 [\text{Cr (VI)}] \quad \dots (3)$$

### Dependence of rate on the concentration of $\alpha$ -hydroxy acid in Polysorbate 80 and DDAC:

The oxidation studies were carried out by varying initial [ $\alpha$ -hydroxy acid] in the range 1.00 to 4.00  $\times 10^{-3}$  mol dm<sup>-3</sup> by keeping other variables constant. The near consistency in the k<sub>2</sub> values (Tables 4 and 5 ) and the slope of nearly unity is obtained from a linear graph of logarithm of specific rate ( k<sub>1</sub> in s-1 ) verses logarithm of  $\alpha$ -hydroxy acid concentration in each case suggesting first order dependence of rate on [ $\alpha$ -hydroxy acid] (Figs. 3 and 4). Hence the rate law for the Cr (VI) oxidation  $\alpha$ -hydroxy acid of is given below equation 4.

$$-d [\text{Cr (VI)}] / dt = k_2 [\text{Cr (VI)}] [\alpha\text{-hydroxy acid}] \quad \dots (4)$$

### Comparison of rates on oxidation of Pentaammine cobalt (III) complexes of both bound and unbound $\alpha$ -hydroxy acid by QCC:

Specific rate of the lactato complex is more compared to both the rates of oxidation of unbound ligand and Mandela to complex deserves an explanation. The ligation of lactic acid to Co (III) centre has probably increased its reactivity towards QCC and this effect seems to be more specific for this ligand only. If the reaction proceeds through a preformed Chromate ester, then the rate of C-H fission will have been enhanced, resulting in an increased rate of oxidation of lactato complex such a precursor complex may be satirically hindered in the case of Mandela to and glycolato complexes.

### 4. Mechanism

Oxidation of Pentaamminecobalt (III) complexes of both bound and unbound ligands in surfactant medium. Thus, the kinetics of two electron transfer route seems to be unavailable for QCC with Cobalt (III) bound and unbound complexes of  $\alpha$ -hydroxy acid in surfactant medium, QCC oxidizes Cobalt (III) bound and unbound  $\alpha$ -hydroxy acids. It rules out the synchronous C-C bond fission and electron transfer to Cobalt (III) centre. Oxidation of above complexes increase with increase of temperature. With increase in surfactant concentration an increase in the rate is observed. The added DDAC enhances the rate of oxidation of a reaction much more than Polysorbate 80. Similar trends has been observed in lactato and glycolato Co (III) complexes.

Table 1

|  |  |
|--|--|
| [(NH <sub>3</sub> ) <sub>5</sub> Co <sup>III</sup> -L] <sup>2+</sup> | = 2.00 $\times 10^{-2}$ mol dm <sup>-3</sup> |
| [QCC]  | = 2.00 $\times 10^{-2}$ mol dm <sup>-3</sup> |
| [HClO <sub>4</sub> ]   | = 1.00 mol dm <sup>-3</sup>                  |
| [Surfactants]  | = 1.00 $\times 10^{-4}$ mol dm <sup>-3</sup> |
| Temperature  | = 34 $\pm$ 0.2°C                             |

| Time (S) | 10 <sup>-3</sup> (a-x) mol dm <sup>-3</sup> |      |
|----------|---|------|
|          | Polysorbate 80                              | DDAC |
| 180      | 9.50  | 4.10 |
| 360      | 6.66  | 2.72 |
| 540      | 5.14  | 2.07 |
| 720      | 3.69  | 1.50 |
| 900      | 2.72  | 0.96 |
| 1080     | 1.99  | 1.17 |
| 1260     | 1.28  | 0.51 |
| 1440     | 1.07  | 0.37 |
| 1620     | 1.01  | 0.32 |
| 1800     | 0.91  | 0.23 |




Fig 1: First order dependence plots

Table 2

|                      |   |
|----------------------|---|
| [QCC]                | $= 2.00 \times 10^{-3} \text{ mol dm}^{-3}$ |
| [HClO <sub>4</sub> ] | $= 1.00 \text{ mol dm}^{-3}$                |
| [Polysorbate 80]     | $= 1.00 \times 10^{-4} \text{ mol dm}^{-3}$ |
| Temperature          | $= 34 \pm 0.2^\circ\text{C}$                |

|              | $[(\text{NH}_3)_5\text{Co}^{\text{III}}-\text{L}] \text{ } 10^2 \text{ mol dm}^{-3}$ | $10^4 k_1 (\text{S}^{-1})$ | $10^2 k_2 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ |
|--------------|--|----------------------------|---|
| L= Lactato   | 1.00   | 2.96                       | 2.96  |
|              | 2.00   | 3.90                       | 2.95  |
|              | 3.00   | 5.70                       | 2.96  |
|              | 4.00   | 7.70                       | 2.95  |
|              | 5.00   | 10.45                      | 2.95  |
| L= Mandelato | 1.00   | 2.17                       | 2.170   |
|              | 2.00   | 2.85                       | 2.169   |
|              | 3.00   | 3.84                       | 2.164   |
|              | 4.00   | 5.42                       | 2.168   |
|              | 5.00   | 7.26                       | 2.169   |
| L=Glycolato  | 1.00   | 1.65                       | 0.364   |
|              | 2.00   | 1.98                       | 0.368   |
|              | 3.00   | 2.55                       | 0.372   |
|              | 4.00   | 3.41                       | 0.352   |
|              | 5.00   | 4.32                       | 0.366   |


Fig 2: Dependence of rate on  $[\text{Co}^{\text{III}}]$  in Polysorbate 80

Table 3

|                      |   |
|----------------------|---|
| [QCC]                | $= 2.00 \times 10^{-3} \text{ mol dm}^{-3}$ |
| [HClO <sub>4</sub> ] | $= 1.00 \text{ mol dm}^{-3}$                |
| [DDAC]               | $= 1.00 \times 10^{-4} \text{ mol dm}^{-3}$ |
| Temperature          | $= 34 \pm 0.2^\circ\text{C}$                |

|            | $[(\text{NH}_3)_5\text{Co}^{\text{III}}-\text{L}] \text{ } 10^2 \text{ mol dm}^{-3}$ | $10^4 k_1 (\text{S}^{-1})$ | $k_2 \text{ } 10^2 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ |
|------------|--|----------------------------|--|
| L= Lactato | 1.00   | 1.00                       | 1.00   |
|            | 2.00   | 1.62                       | 1.02   |
|            | 3.00   | 2.43                       | 1.02   |
|            | 4.00   | 3.80                       | 1.00   |
|            | 5.00   | 6.10                       | 1.01   |

|              |      |      |       |
|--------------|------|------|-------|
| L= Mandelato | 1.00 | 0.64 | 0.64  |
|              | 2.00 | 0.96 | 0.64  |
|              | 3.00 | 1.55 | 0.64  |
|              | 4.00 | 2.25 | 0.64  |
|              | 5.00 | 3.44 | 0.64  |
| L=Glycolato  | 1.00 | 0.41 | 0.239 |
|              | 2.00 | 0.67 | 0.238 |
|              | 3.00 | 0.96 | 0.237 |
|              | 4.00 | 1.59 | 0.239 |
|              | 5.00 | 2.39 | 0.238 |




Fig 3: Dependence of rate on [Co<sup>III</sup>] in DDAC

Table 4

[QCC] = 2.00× 10<sup>-3</sup> mol dm<sup>-3</sup>  
[HClO<sub>4</sub>] = 1.00 mol dm<sup>-3</sup>  
[Polysorbate 80] = 1.00× 10<sup>-4</sup> mol dm<sup>-3</sup>  
Temperature = 34±0.2°.

| [α – Hydroxy acid]10 <sup>2</sup> mol dm <sup>-3</sup> | 10 <sup>4</sup> k <sub>1</sub> (S <sup>-1</sup> ) | 10 <sup>2</sup> k <sub>2</sub> dm <sup>3</sup> mol <sup>-1</sup> s <sup>-1</sup> |
|--|---|--|
| L= Lactic acid   | 1.00  | 1.23   |
|  | 2.00  | 2.01   |
|  | 3.00  | 3.19   |
|  | 4.00  | 5.12   |
|  | 5.00  | 8.01   |
| L= Mandelic acid                                       | 1.00  | 0.83   |
|  | 2.00  | 1.28   |
|  | 3.00  | 2.00   |
|  | 4.00  | 3.02   |
|  | 5.00  | 5.14   |
| L=Glycolic acid  | 1.00  | 0.45   |
|  | 2.00  | 0.74   |
|  | 3.00  | 1.01   |
|  | 4.00  | 1.72   |
|  | 5.00  | 2.53   |





Fig 4: Dependence of rate on [α – hydroxy acid] in Polysorbate 80

Table 5

|                      |   |
|----------------------|---|
| [QCC]                | $= 2.00 \times 10^{-3} \text{ mol dm}^{-3}$ |
| [HClO <sub>4</sub> ] | $= 1.00 \text{ mol dm}^{-3}$                |
| [DDAC]               | $= 1.00 \times 10^{-4} \text{ mol dm}^{-3}$ |
| Temperature          | $= 34 \pm 0.2^\circ$                        |

| $[\alpha\text{-hydroxy acids}] 10^2 \text{ mol dm}^{-3}$ | $10^4 k_1 (\text{S}^{-1})$ | $10^2 k_2 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ |
|--|----------------------------|---|
| Lactic acid  | 1.00                       | 1.29  |
|  | 2.00                       | 2.24  |
|  | 3.00                       | 3.88  |
|  | 4.00                       | 6.32  |
|  | 5.00                       | 11.80   |
| Mandelic acid  | 1.00                       | 0.65  |
|  | 2.00                       | 1.04  |
|  | 3.00                       | 1.60  |
|  | 4.00                       | 2.54  |
|  | 5.00                       | 3.99  |
| Glycolic acid  | 1.00                       | 0.41  |
|  | 2.00                       | 0.67  |
|  | 3.00                       | 1.16  |
|  | 4.00                       | 1.85  |
|  | 5.00                       | 3.24  |

Fig 5: Dependence of rate on  $[\alpha\text{-Hydroxy acid}]$  in DDAC

## 5. Conclusion

The oxidation reactions of Cobalt (III) complexes of  $\alpha$ -hydroxy acids viz, lactic acid, glycolic acid and mandelic acid have been carried out using a novel chromium (VI) oxidant. There is a remarkable increase in the rate of the reaction has been observed in the presence of novel surfactants Polysorbate 80 and DDAC. These surfactants act as a positive catalyst in situ of the reaction and among which DDAC enhances the rate much more than the Polysorbate 80.

## 6. Acknowledgments

Authors are thanks to the Management of Vaniyambadi Muslim Educational Society and PG& Research Department of Chemistry, Islamiah College Vaniyambadi for providing the necessary facilities for the successful completion of this research work. At last I thanks to my friend R. Kalaivani M.A., for his kind help in processing the complete this work in computer.

## 7. References

- Kresheilk GC, in water Comprehensive Treatment CF. Frank Ed Plenum. 1973; 4:23.
- Bunton CA, Kamega A, Mich K. J org Chem. 1972; 37:1388.
- Bunton CA. In Reaction Kinetics micelles. E.H.Cordese Plenum. 1972; 73:1388.
- Zaffren E, Watson R E. Intrsci Che Rep, 1972, 6: 51.
- Gurumurthy R, Gopalakrishnan M, Karthikeyan B, Selvaraju M. Asian J Chem. 1998; 10(3):476.
- Saran NK, Acharya RC. Indian Chem Soc. 1985; 62:747.
- Haque ME, Das AR, Rakshit AK, Moulik SP. Properties of Mixed Micelles of Binary Surfactant Combinations. Langmuir. 1996; 12:4048-4089.
- Brahmaiah A, Manikyamba P. Indian J Chem. 1995; 34(2):900.
- BalojKawle, ThirupathiRao M, Adinarayan M. India J Chem. 1994; 33(2):1021.
- Balasubramanian K, Lakshmanan K, Sekar G. Asian J Chem. 1999; 11:1451.
- Sankaran KR, Anbuselvan C. Asian J. Chem. 1998; 10(4):806.
- Ramkumar B. Afinidad. 2003; 60(505):257-261.
- Mohmed Farook A. J Sol. Chem. 2007; 36:345-356.
- Subramian K, Udhayavani S. Journal of Chem. Bio and Phy sciences. 2012; 2(2):551-559.
- Anandaratchagan B, Md.Nawaz, Subramani K. Acta. Chim. Pharm India. 2011; 1(1):44-50.

16. Thaminum Ansari A. Int.Journal of Chem Tech Research. 2009; 1(2):308-313.
17. Siebert H, Anorg Z, Allg Chem. 1978; 47:441.
18. Kalidoss P, Srinivasan VS, Indian J Chem. 1987; 26:924-929.
19. Kalidoss P, Srinivasan VS. Indian J Chem. 1987; 26:211-214.
20. Kalyan Kalisen Gupta, Amalendu Banerjee, Hrishikesh Chatterjee, Tetrahedron. 1992; 48, 25(30):5323.
21. Abdul Vaheith Z, Mansur Ahmed. A Kinetic Study in Oxidation of Cobalt (III). Complexes of Alpha-Hydroxy Acids by Quinolonium Bromo Chromate in Na LS. Medium, International Journal of Frontiers in Science and Technology. 2015; 3(3):49-58.
22. Sekar KG, Sakthivel RV. Reactivity of Cyclanols towards Quinaldinium Fluorochromate Oxidation, J Solution Chem. 2013; 42:1748-1756.