



## Isolation and Partial Characterization of a Flavanone from Sudanese *Mitragyna inermis* (Willd.) O. Kuntze (Rubiaceae) Stem Bark

Abdel Karim M<sup>1\*</sup>, Waladeen S<sup>2</sup>, Inas OK<sup>3</sup>

<sup>1-2</sup> Sudan University of Science and Technology, Faculty of Science, Sudan

<sup>3</sup> University of Bahri, College of Applied and Industrial Sciences, Sudan

### Abstract

Genus *Mitragyna* is distributed in tropical and subtropical regions of Africa and Asia. Different species of *Mitragyna* are widely used in ethnomedicine against malaria, fever, worms, diarrhea and cough. *Mitragyna speciosa* is a natural remedy for fatigue. *Mitragyna ciliate*, *Mitragyna inermis* and *Mitragyna stipulosa* have been used traditionally against hypertension, inflammation, rheumatism, gonorrhoea and bronchopulmonary diseases. *Mitragyna africanus* is used traditionally against mental diseases. In this study stem barks of *Mitragyna inermis* were extracted with 95% ethanol and the crude extract was fractionated over silica gel plates to yield a chromatographically pure flavonoid – compound I. The structure of the isolated compound has been partially characterized by its spectral data (UV and <sup>1</sup>HNMR).

**Keywords:** *Mitragyna inermis*, flavonoid, isolation, partial characterization

### 1. Introduction

*Mitragyna* is a genus in the family Rubiaceae. This genus is distributed in tropical and subtropical regions of Africa and Asia. The genus contains ten species: *Mitragyna parvifolia*, *Mitragyna hirsuta*, *Mitragyna diversifolia*, *Mitragyna tubulosa*, *Mitragyna speciosa*, *Mitragyna rotundifolia*, *Mitragyna ciliate*, *Mitragyna inermis*, *Mitragyna africanus* and *Mitragyna stipulosa* [1].

Different species of *Mitragyna* are widely used in ethnomedicine [2, 3], against malaria, fever, worms, diarrhea and cough. *Mitragyna speciosa* is a natural remedy for fatigue [4]. *Mitragyna ciliate*, *Mitragyna inermis* and *Mitragyna stipulosa* have been used traditionally against hypertension, inflammation, rheumatism, gonorrhoea and bronchopulmonary diseases. *Mitragyna africanus* is used traditionally against mental diseases [5].

Some indole alkaloids, triterpenoids and saponins have been reported from the genus *Mitragyna*. The major alkaloid of this genus demonstrated a wide reaching pharmacological potential including cardiovascular, antitumor, and antimicrobial activities [6-9]. It has been documented that mitragynine-the major alkaloid of *Mitragyna*- possesses analgesic properties [10-12]. Different extracts of *Mitragyna speciosa* and *Mitragyna parvifolia* showed significant antiinflammatory activity [13-16]. It has been reported that *Mitragyna speciosa* exhibited significant antimutagenic activity [17]. *Mitragyna ciliate* extracts induced aortic relaxation in a dose-dependant manner [18]. In some in vivo studies, The extracts of *Mitragyna diversifolia* exhibited antidiarrheal effect [19]. Extracts of *Mitragyna speciosa* also showed antidiarrheal activity [20, 21]. The antimicrobial and antioxidant activities of *Mitragyna speciosa* have been reported [22].

### 2. Materials and Methods

#### 2.1 Materials

##### 2.1.1 Plant material

*Mitragyna inermis* barks were collected from Damazin (Sudan). The plant was identified and authenticated by the Medicinal and Aromatic Plants Research Institute, Khartoum – Sudan.

#### Instruments

UV spectra were run on a Shimadzu 2401PC UV- Visible Spectrophotometer. <sup>1</sup>HNMR spectra were performed on a Joel ECA 500MHZ NMR Spectrophotometer.

#### Methods

##### Extraction and Isolation of flavonoid

Air dried powdered *Mitragyna inermis* barks (1 kg) were macerated at room temperature with ethanol (95%) for 72h. The solvent was evaporated under reduced pressure to dryness to give a crude product.

The crude extract was fractionated via thin layer chromatography using 30% acetic acid as solvent. The chromatograms were viewed and located under UV light and a flavonoid-compound I- was eluted from silica with methanol.

#### Results and Discussion

*Mitragyna inermis* barks crude extract was fractionated over silica gel plates to yield a chromatographically pure flavonoid – compound I. The structure of the isolated compound has been partially characterized via some spectral data (UV and <sup>1</sup>HNMR).

### Characterization of compound I

The UV spectroscopy is a valuable analytical tool in the chemistry of flavonoids and can differentiate between flavonoids with double bond in position 2-3 and those lacking such bond. The UV spectra of most flavonoids consist of two major absorption maxima one of which occur in range 220-285 nm (band II) and the other in the range 300-400nm (band I) the appearance of both band I and II in the UV spectrum demonstrates conjunction between the benzoyl and cinnamoyl system.




Fig 1

The UV spectrum of compound I (Fig. 3) showed  $\lambda_{max}$  (MeOH) 249m. The appearance of only one band – band II- in this spectrum suggests saturation at the C<sub>2</sub> – C<sub>3</sub> position. Consequently compound I is probably a (i) flavanone, (ii) isoflavone (iii) dihydrochalcone or (iv) a dihydroflavonol.




Fig 2

However, isoflavones are easily distinguished by a shoulder in the UV range 300-340nm. Such shoulder has not been detected in the UV spectrum of compound I. On the other hand dihydroflavonols are characterized by a 3-OH function which could be detected by the UV shift reagent- sodium methoxide. Sodium methoxide can give a bathochromic shift with decrease in intensity in presence of a 3-OH. The sodium methoxide spectrum (Fig. 4) failed to give a bathochromic shift.




Fig 1: UV spectrum of compound I




Fig 2: Sodium methoxide spectrum of compound I




Fig 3: Sodium acetate spectrum




Fig 4: Aluminium chloride spectrum of compound I




Fig 5: Boric acid spectrum of compound I

In their  $^1\text{H}$ NMR spectra flavanones show a double multiplet around 2.8ppm and 5.2ppm. The mutual spin-spin splitting of the magnetically nonequivalent protons at  $\text{C}_3$  suffers further splitting by the neighboring  $\text{C}_2$  protons to give a double doublet (usually merging into a multiplet) around 2.8ppm. The signal of  $\text{C}_2$  proton is split by one of the protons at  $\text{C}_3$  into a doublet. Such doublet suffers further splitting by the other proton at  $\text{C}_3$  to yield a double doublet (usually appearing a multiplet) around 5.2ppm. However, these multiplets were detected in the  $^1\text{H}$ NMR spectrum of compound I (Fig. 8). Thus the isolated flavonoid is a flavanone.

The hydroxylation pattern of compound I has been studied via various UV shift reagents: sodium acetate (which gives a bathochromic shift in presence of a 7-OH); aluminium chloride (showing a bathochromic shift diagnostic of 3-, 5-OH and catechol systems) and boric acid (diagnostic of catechol moieties).

When a methanolic solution of compound I was treated with the shift reagent: sodium acetate, no bathochromic shift was observed indicating absence 7-hydroxylation (Fig. 5).

The aluminium chloride spectrum failed to show a bathochromic shift. This clearly suggests absence of 3-, 5-OH as well as catechol systems (Fig. 6). The boric acid spectrum behaved in the same manner and did not show any bathochromic shift suggesting absence of catechol systems (Fig. 7).

The  $^1\text{H}$ NMR spectrum showed  $\delta(\text{ppm})$ : 1.25, 1.52 (assigned for two methyl groups), multiplet centered at 2.00 (assigned for  $\text{C}_3$  protons), multiplet: 3.60-4.50 (sugar protons-not identified in this study), multiplet centered at 5.20 (accounting for  $\text{C}_2$  proton). The aromatic protons appeared as a multiplet centered at 7.40ppm. Signals at 2.50 and 3.35 are due to solvent (DMSO) residual protons and residual water respectively.





Fig 6:  $^1\text{H}$ NMR spectrum of compound I

On the basis of the above argument, the following partial structure was proposed for compound I:



## References

1. Shellard EJ, Houghton PJ, Payom T, Masechaba R, Phillipson JD. The *Mitragyna* species of Asia part XXX. Journal of Medicinal Plant Research. 1978; 34:253.
2. Shellard EJ, Phillipson JD. The *Mitragyna* species of Asia. Part I. The alkaloids of the leaves of *Mitragyna rotundifolia* (Roxb.) O. Kuntze. Journal of Medicinal Plant Research. 1964; 12:27-32.
3. Shellard EJ, Houghton PJ. The distribution of alkaloids in *Mitragyna parvifolia* (Roxb.) Korth in young plants grown from Ceylon seed. Journal of Pharmacy and Pharmacology. 1971; 23:245.
4. Sangun Suwanlert. A Study of Kratom Eaters in Thailand. Bulletin on Narcotics. 1975; 27:21-27.
5. Moklas MAM, Nurul Raudzah AR, Taufik Hidayat M, Sharida F, Farah Idayu N, Zulkhairi A *et al.* A Preliminary Toxicity Study of Mitragynine, An Alkaloid from *Mitragyna speciosa* Korth and its Effects on Locomotor Activity in Rats. Advances in Medical and Dental Sciences. 2008; 2:56-60.
6. Shellard EJ, Becket AH, Payom T *et al.* The *Mitragyna* species of Asia part V. Journal of Medicinal Plant Research. 1967; 15:245-254.
7. Shellard EJ, Houghton PJ. The *Mitragyna* species of Asia part XXV. Journal of Medicinal Plant Research. 1974; 25:172-174.
8. Takayama H. Chemistry and pharmacology of analgesic indole alkaloids from the Rubiaceae plant, *Mitragyna speciosa*. Chemical & Pharmaceutical Bulletin. 2004; 52:916-28.
9. Kang WY, Hao XJ. Triterpenoids from *Mitragyna rountifolia*. Biochemical Systematics and Ecology. 2006; 34:585-587.
10. Watanabe K, Yano S, Horie S, Yamamoto LT. Inhibitory effect of mitragynine, an alkaloid with analgesic effect from Thai medicinal plant *Mitragyna speciosa*, on electrically stimulated contraction of isolated guinea-pig ileum through the opioid receptor. Life Sciences. 1997; 60:933-942.
11. Yamamoto LT, Horie S, Takayama H, Aimi N, Sakai S, Yano S *et al.* Opioid receptor agonistic characteristics of mitragynine pseudoinoxyl in comparison with mitragynine derived from Thai medicinal plant *Mitragyna speciosa*. General Pharmacology. 1999; 33:73-81.
12. Takayama H, Ishikawa H, Kurihara M, Kitajima M, Aimi N, Ponglux D, *et al.* Studies on the synthesis and opioid agonistic activities of mitragynine-related indole alkaloids: discovery of opioid agonistic structurally different from other opioid ligands. Journal of

- Medicinal Chemistry. 2002; 45:1949-1956.
13. Winter CA, Risley EA, Nuss GW. Carrageenan-induced edema in hind paw of the rat as an assay for the anti-inflammatory drugs. Proceedings of the Society for experimental Biology and Medicine. 1962; 111:544-547.
  14. Dongmo AB, Kamanyi A, Dzikouk G, Chungag-Anye Nkeh B, Tan PV, Nguelefack T, *et al.* Anti-inflammatory and analgesic properties of the stem bark extract of *Mitragyna ciliata* (Rubiaceae) Aubre´v. and Pellegr. Journal of Ethnopharmacology. 2003; 84:17-21.
  15. Gupta V, Kumar P, Bansal P, Singh R. Anti-inflammatory and Anti-nociceptive Activity of *Mitragyna parvifolia*. Asian Journal of Medical Sciences. 2009; 1:97-99.
  16. Shaik Mossadeq WM, Sulaiman MR, Tengku Mohamad TA, Chiong HS, Zakaria, ZA, Jabit ML, *et al.* Anti-Inflammatory and Antinociceptive Effects of *Mitragyna speciosa* Korth Methanolic Extract. Medical Principles and Practice. 2009; 18:378-384.
  17. Ghazali AR, Abdullah R, Ramli N, Rajab NF, Ahmad-Kamal M, Yahya NA. Mutagenic and antimutagenic activities of *Mitragyna speciosa* korth extract using Ames test. Journal of Medicinal Plants Research. 2011; 5:1345-1348.
  18. Dongmo1 A, Kamanyi MA, Tan PV, Bopelet M, Vierling W, Wagner H. Vasodilating Properties of the Stem Bark Extract of *Mitragyna ciliata* in Rats and Guinea Pigs. Phytotherapy Research. 2004; 18:36-39.
  19. Jebunnessal Uddin SB, Mahabub-Uz-Zaman M, Akter R, Ahmed NU. Antidiarrheal activity of ethanolic bark extract of *Mitragyna diversifolia*. Bangladesh Journal of Pharmacology. 2009; 4:144-146.
  20. Chittrakarn S, Sawangjaroen K, Prasettho S, Janchawee B, Keawpradub N. Inhibitory effects of kratom leaf extract (*Mitragyna speciosa* Korth.) on the rat gastrointestinal tract. Journal of Ethnopharmacology. 2008; 116:173-178.
  21. Matsumoto K, Hatori Y, Murayam T, Tashima K, Wongseripatana S, Misawa K *et al.* Involvement of  $\mu$ -opioid receptors in antinociception and inhibition of gastrointestinal transit induced by 7-hydroxymitragynine, isolated from Thai herbal medicine *Mitragyna speciosa*. European Journal of Pharmacology. 2006; 549:63-70.
  22. Parthasarathy S, Azizi JB, Ramanathan S, Ismail S, Sasidharan S, Mohd MI, Said, i Mansor SM. Evaluation of antioxidant and antibacterial activities of aqueous, methanolic and alkaloid extracts from *Mitragyna speciosa* (Rubiaceae family) leaves. Molecules. 2009; 14:3964-3974.